

TECHNICAL DATA SHEET

Article No. 9166

MYP Agar, prepared plates

SYNONYMS

Mannitol Egg Yolk Polymyxin Agar, Bacillus Cereus Mossel Agar

SPECIFICATION

Prepared plates. Selective solid medium, according to Mossel, for the isolation and identification of Bacillus cereus from food samples. FIL-IDF 181, ISO 7932, ISO 21871.

Color: Orange
pH: 7.2 ± 0.2 at 25 °C

COMPOSITION IN G/L

Casein peptone	10.000
Mannitol	10.000
Sodium chloride	10.000
Meat extract	1.000
Phenol red	0.025
Agar	12.000
Polymixin B Sulphate	100,000IU
Egg Yolk	100 ml

PACKAGING DETAILS

9166-20PLATES

20 prepared plates 90 mm

Content: 21 ± 2 ml

Packaging unit: 1 box with 2 packs of 10 plates/pack. Single cellophane.



GUIDELINES

Description:

Mossel's formulation has been developed to detect and enumerate *B. cereus* in any food stuff. It is both selective and differential for this microorganism. Polymyxin addition inhibits most of the accompanying bacteria, but does not affect the growth of *B. cereus*. This bacterium does not ferment mannitol and thus there is no change in the colour of the indicator around the colonies. The lecithinase activity of *B. cereus* produces a halo or zone of white precipitate around the colonies.

A count of *B. cereus* of more than 100.000 cells/g of food sample is considered hazardous, since the accumulated phosphorilcholine may cause toxic symptoms in children. For this reason a viable enumeration has to be performed to evaluate the actual population of cells.

Technique:

For plate inoculation follow the laboratories' standard methods or the applicable norms (spiral plating method, econometric methods, streak plating, dilution banks, spread plating with drigralsky rod, among them).

According to the authors, dehydrated or dry samples must be treated in the following way: 20 g of sample is mixed with 90 ml of Tryptone Water for a minimum period of 1 hour, at room temperature. Afterwards, add an additional 90 mL of Tryptone Water and homogenize. If necessary dilute 1:10. Proceed to a 1/10 serial dilution bank using Tryptone water as the diluent if necessary. With a Drigalsky loop, spread aliquots of 0,1 mL over the surface of the agar plates and let the agar medium absorb the aliquots. Incubate the plates at 30 °C for 18-24 hours to allow spore germination before giving definite results.

Suspected colonies have the following appearance: irregular borders, pink colour becoming purple in the centre, with a halo of white precipitate (mannitol +). Colonies with yellow halos must be discounted (mannitol -).

Confusion with other colonies of Gram positive bacilli is possible, and hence, confirmation tests must be carried out i.e. glucose fermentation, gelatine degradation and nitrate reduction.

MICROBIOLOGICAL CONTROL

Inoculate: Practical range 100 ± 20 CFU. min. 50 CFU (productivity)/ 10⁴-10⁶ (selectivity).

Microbiological control according to ISO 11133:2014/A1:2018.

Aerobiosis. Incubation at 30 ± 1 °C, read after 24 ± 3h - 44 ± 4h.

Microorganism	Growth
<i>Bacillus cereus</i> ATCC® 11778, WDCM 00001	Good-pink colonies with halo of precipitation
<i>Escherichia coli</i> ATCC® 25922, WDCM 00013	Inhibited
<i>Bacillus subtilis</i> ATCC® 6633, WDCM 00003	Yellow colonies without halo

Sterility control:

Incubation 48 hours at 30-35 °C and 48 hours at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.



BIBLIOGRAPHY

- ATLAS, R.M. & L.C. PARKS (1993) Handbook of Microbiological Media. CRC Press. London.
- CORRY, J.E.L., G.D.W. CURTIS & R.M. BAIRD. (2003) Handbook of Culture Media for Food Microbiology. Elsevier Sci. B.V. Amsterdam. The Netherlands.
- DOWNES, F.P. & K. ITO (2001) Compendium of methods for the microbiological examination of foods. 4th ed. APHA. Washington DC. USA.
- FIL-IDF 181:1998 Provisional Int. Standard. Dried Milk Products. Enumeration of Bacillus cereus.- Most probable number technique.
- ISO 7932 Standard (2004) 3rd ed. Microbiology of food and animal feeding stuffs. Horizontal method for the enumeration of presumptive Bacillus cereus. Colony count technique at 30°C.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- ISO 21871 Standard (2006) Microbiology of food and animal feeding stuffs.- Horizontal method for the determination of low numbers of presumptive Bacillus cereus.- Most probable number technique and detection method.
- MOSSEL, D.A.A., KOOPMAN. M.J. & JONGERIUS, E. (1967) Enumeration of Bacillus cereus in foods. Appl. Microbiol. 15:650-653.
- PASCUAL ANDERSON, M^a.R^a (1992) Microbiología Alimentaria. Díaz de Santos, S.A. Madrid

STORAGE

2-14 °C

SHELF LIFE

3 months unopened from date of manufacture

updated: 30.08.2022

